

Hand Infections

In the management of hand infection take care:

- 1- General and local rest to the hand in the elevated position.
- 2- Early administration of antibiotics.
- 3- Early drainage of infection i.e. *don't wait for fluctuation*.
- 4- Incision must be planned and under general or regional anesthesia.
- 5- Incision is done under tourniquet to have a dry field to be able to explore the abscess.
- 6- The hand must be maintained in the elevated position to prevent edema & congestion.
- 7- Position of immobilization is that of function of the hand.
- 8- Early restoration of function by movement.

بدایة كلامك في أي سؤال

Pulp Space Infection

Treatment:

- 1- Don't wait for fluctuation.
- 2- The incision is sited directly over the most tender point and in direction of Langer's lines.
- 3- Other technique: 2 incisions on sides of the palm to divide all septa + drain.


Web Space Infection

Management:

- 1- A transverse skin incision is made over the point of maximum tenderness or fluctuation and then an artery or a sinus forceps is opened in a longitudinal direction (to avoid damage to the digital nerves and vessels), through the subcutaneous tissue to enter the abscess cavity.
- 2- Vertical dorsal incision. Take care, and don't reach the edge of the web.

Hilton

Facial Spaces



Hypothenar Space Infection

Management

- A vertical incision at the site of maximal tenderness or along the medial border of the fifth metacarpal bone is done. The space is entered by a sinus forceps i.e. **Hilton's method**.

Thenar Space Infection

Management

- Transverse incision at the web then the space is open by a sinus forceps (Hilton's method) **OR**
- Vertical incision on the lateral aspect of the back of the 2nd metacarpal bone, this is the commonly used incision.


Mid Palmar Space Infection

Management

- Incision when pus is formed: either:

- If there is a collar-stud abscess, incise along one of the hand creases, draining the subcutaneous abscess. The hole in the aponeurosis is explored and is enlarged to drain subaponeurotic collection.
- A transverse incision is done like that of the web space and by Hilton's method a sinus forceps is introduced into the retrotendinous space to drain.

Incisions Of the Hand



Jaw Swellings	
<p>III) Bone Swellings</p> <p>as - Ivory osteoma</p> <p>- Osteo clastoma</p> <p>-Maxillary antrum carcinoma</p> <p>- Fibrosarcoma</p> <p>- Osteosarcoma of the mandible</p>	<p>Giant cell granuloma (Benign)</p> <p>osteoclastoma (malignant)</p> <p>- Bone tumour</p> <p>- Synthesis menti</p> <p>- Outer & inner tables</p> <p>- see Later</p> <p>- see Later</p> <p>- same</p> <p>- Unequal cysts → Soap bubble appearance</p> <p>- Same + other option → Curettage & Radiotherapy</p>
<p>III) Odontomes</p> <p>• May arise from ectodermal or mesodermal origin of tooth or both.</p> <p>• Those of ectodermal origin (Paradental debris of Malassez) are common.</p>	<p>III) Odontomes</p> <p>• May arise from ectodermal or mesodermal origin of tooth or both.</p> <p>• Those of ectodermal origin (Paradental debris of Malassez) are common.</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p> <p>- Younger age</p> <p>- Wall → same content is clear & glory + contains tooth</p> <p>- same</p> <p>- Cystic swelling with tooth inside</p> <p>- Same</p> <p>- Derroofing of the cyst & the lining membrane.</p> <p>Tooth → Normally directed → Not removed</p> <p>Maldirected → Removed</p>
<p>A) Dental Cyst</p> <p>• origin</p> <p>• etiology</p> <p>• site</p> <p>• age</p> <p>• structure</p> <p>• C/P</p> <p>• X-ray</p> <p>• D.D.</p> <p>• treat.</p>	<p>B) Dentigerous cyst</p> <p>- same</p> <p>- Unrupted tooth</p> <p>- Lower 3rd molar</p>

Tongue Ulcers

AH

Ahmed El.Sherbiny 13-12-2004


East or West	Inflammatory			Neoplastic		
	Traumatic	Acute	Chronic	(Squamous Cell Carcinoma)		
AH	Dental Ulcer	Dyspeptic (Apthus)	TB	Syphilitic		
Is The Best	Frenular Ulcer					
Etiology	Sharp teeth	Unknown (was wrongly thought upper GIT disturb.)	2ry to open Pulmonary TB	Breakdown of gumma	- Previous irradiation - Marjolin's ulcer - Carcinogenic agents	
Number	Single	Multiple	May be multiple	Single	Single	
Pain	Painful	Painful	Severe	Painless	Pain related to the ear	
Site	Margin of the tongue	Sides, dorsum & inner surface of lip & cheek	Dorsum posteriorly & tip anteriorly	Dorsum in midline	- Lat. margin of ant. 2/3 50% - Posterior 1/3 20%	
Size	Usually small	Small	Usually small	Large	Grow rapidly & may reach large size	
Shape	Rounded	Rounded or oval	Rounded	Rounded	Irregular	
Surface	Granulation tissue	Covered with white scabs	Pale granulation tissue	Wash-leather (dirty)	Malignant fungating tissue	
Margin	Hyperemic	Hyperemic	Cyanotic		Necrotic tissue	
Edge	Sloping	Sloping	Undermined & cyanotic	Punched out	Raised everted	
Base	Soft	Soft	Soft	Indurated	Indurated	
L.N.	Chronic non-specific	Chronic non-specific	TB lymphadenitis		Metastatic LNs: Hard, progressive and fixed	
TTT	Remove septic tooth	- Gentian violet - Anesthetic gel.	- Anti TB ttt - Oral hygiene	Anti syphilitic drugs	- Surgical removal - Radiotherapy	

NB. 1- Chronic Superficial Glossitis : see AH notes

2- Acute inflammatory ulcers also includes:

- Herpetic ulcer which is multiple, painless, rounded, with hyperemic margin and is caused by herpes simplex virus. It is treated with antiviral drugs.
- Lichen planus which is supposed to be due to an autoimmune mechanism. It affects the skin, oral mucosa and may cause ulcers + hyperkeratotic lesions




TREATMENT OF CHRONIC ISCHEMIA

	CONSERVATIVE METHODS	INTERVENTIONAL RADIOLOGICAL PROCEDURES	SURGICAL TREATMENT		
			Direct arterial surgery (Arterial reconstruction)	Indirect arterial surgery (Lumbar sympathectomy)	Amputation
Indications	1- Mild ischemia. 2- Poor general condition of the patient. 3- When an operation is not technically feasible	Localized obstruction in large and medium sized arteries.	a- Severe ischemia (rest pain or pre-gangrene). b- Adequate run off. c- Proximal arterial occlusion (anywhere from abdominal aorta to the terminal part of popliteal artery). d- Good general condition	Indications: Cases of ischemia when direct arterial surgery is NOT feasible i.e. with distal occlusion (Bad distal run-off). It is of value in cases: a- With ischemic ulcers b- With rest pain. Contraindications: 1- Intermittent claudication (worsen the muscle ischemia). 2- Gangrene (ineffective)	There are 2 possibilities: 1- <u>Conservative Amputation</u> : If blood supply of adjacent tissues is good or can be improved, line of demarcation appears and separation proceeds by aseptic ulceration. In such cases do either: a- Excision of toes at line of demarcation leaving raw surface to heal by granulation. b- Transmetatarsal or midtarsal amputation in forefoot gangrene provided there is viable long plantar flap reaching base of toes, to cover the stump (dorsal skin cannot withstand trauma). 2- <u>Urgent high amputation</u> : Indications: 1) Spreading gangrene endangering the patient's life. 2) Uncontrollable infection and toxemia. 3) Severe pain deteriorating the general condition of the patient. <u>Level of amputation</u> : Depends on the blood supply sufficient for wound healing: a- In atherosclerosis (involve femoral art.) above knee amputation (the stump will be supplied by the profunda femoris). b- If popliteal pulse is felt below knee amputation
Methods	1- Mild exercise short of causing pain (help the collateral circulation). 2- Stop Smoking. 3- Correction of anemia. 4- Control of DM, hypertension and hyperlipidemia. 5- Drugs: ↓↓ platelet aggregation as dipyridamole, small dose of aspirin, pentoxifylline or PGE. 6- <u>Protection of ischemic parts</u> : a) Carefully washed, dried, and powdered. b) Protected by woolen socks (in winter) and suitable shoes. c) Nails and corns are cut cautiously. d) Infections are treated properly.	1- <u>Percutaneous transluminal angioplasty (PTA)</u> : a special balloon catheter is introduced percutaneously until it reaches the stenosed or occluded segment then inflated to dilate the stenosed segment. 2- <u>Application of stent</u> : After balloon angioplasty 3- <u>Destruction of the atheroma</u> by laser can be performed before angioplasty.	A) <u>Thromboendarterectomy</u> : • Indication: Large artery (e.g. aorta and common iliac) and localized lesions. • Removal of the thickened atherosclerotic intima with the overlying thrombus using endarterectomy loop. B) <u>Bypass grafting</u> : • Indication: Big and medium sized artery with extensive or multiple lesions involving a long segment. • Types of arterial grafts: 1- <u>Synthetic grafts</u> : Teflon, Dacron or PTFE. Indication: Large arteries e.g. Aorto-iliac segment. 2- <u>Autogenous grafts</u> : Using long Saphenous either: a. <u>Reversed long Saphenous vein graft</u> : to prevent obstruction by its valves. It is the graft of choice for femoro-popliteal segment. b. <u>In situ long Saphenous vein graft</u> : valves are destroyed using special valve stripper. All tributaries are ligated to prevent A. V. fistula.	 Thromboendarterectomy Bypass grafting	

A.H.

RAMY ALKONAIYESY

- Direct arterial operations are salvage procedures & should not be performed for early mild cases (e.g. intermittent claudication) as if the operation fails loss of the limb may occur.
- Claudication is a relative indication for surgery e.g. if the patient may lose his job.
- In Thromboendarterectomy before closing the arteriotomy incision, the distal intima should be attached to arterial wall by interrupted sutures to prevent its dissection later on.
- If amputation is to be done bilateral, do your best to do below knee on one side.



- 7 or 8 Interossei → Abd. & add. of fingers
- 2 med. Lumbricals → writing → Flexion MCP Extension IP.
- add. pollicis
- 3 hypotenar → Abd. digiti Minimi

- 3 thenar → opp. pollicis
- Flexor P. Brevis
- Abd. P. Brevis

- 2 lat. Lumbricals.

Small area at the base of the thumb.

Pathology

N.B. Any mucinous carcinoma carries bad prognosis except in breast
Any tumor with lymphocytic infiltration carries good prognosis

Yohanned Pafeg

Treatment of Thyrotoxicosis:


- A) Non-specific: 1) Rest. 2) Sedatives. 3) Nutritive diet and excess fluid. (For all cases).
 B) Specific: 1) Antithyroid drugs. 2) Radioactive Iodine. 3) Surgery.

In Specific ttt: (Each patient is considered Separately and one measure or A combination of measures is chosen for him)

Aim	Medical Treatment	Radioactive Iodine	Surgery
Indications 1- To restore pt to an euthyroid status then, 2- Prescribe a maintenance dose for a prolonged period hoping for permanent remission. 1- Iry thyrotoxicosis. 2- Mild thyrotoxicosis. 3- Small gland. 4- Children and young patients. 5- Pre-operative preparation. 6- Post-operative recurrence. 7- Refusal of surgery. 8- Unfit for surgery.	To destroy the thyroid cells thus reducing the mass of functioning thyroid tissue. 1- Diffuse toxic goiter (> 25 years). 2- Thyrocardiac patients. 3- Refusal of surgery. 4- Recurrence after surgery. 5- Toxic nodule.	1- To decrease the mass of overactive tissues. OK 2- To remove All the overactive tissue in case of toxic nodule. 1- 2ry toxic goitre.* (NB: * = Surgery only). 2- Severe 1 ry thyrotoxicosis 3- Retrosternal toxic goiter*. 4- Huge goitre.* 5- Suspicious of malignancy.* 6- Failure of medical treatment. 7- Occurrence of side effects due to medical ttt. 8- Thyrocardiac after proper preoperative prep. 9- Pregnancy and lactation.* 10- Toxic nodule.	1- Mild cases. 2- Young patient (< 25 years). 3- Recurrence after surgery.
Contraindications 1- Toxic nodular goitre. 2- Retrosternal goitre. 3- Huge goitre causing pressure symptoms. 4- Suspicious of malignancy. 5- Leucopenia or agranulocytosis. 6- Pregnancy and lactation.	1- Below the age of 25 years 2- Retrosternal goitre. 3- Huge goitre. 4- Suspicious of malignancy. 5- Pregnancy and lactation. 6- 2ry toxic goitre.	1- Below the age of 25 years 2- Retrosternal goitre. 3- Huge goitre. 4- Suspicious of malignancy. 5- Pregnancy and lactation. 6- 2ry toxic goitre.	1- Mild cases. 2- Young patient (< 25 years). 3- Recurrence after surgery.
Method • <u>Methods:</u> a) <u>Sedatives:</u> Aim: - To insure complete mental rest and good night sleep. b) <u>Indral</u> (β blocker); Action: - \downarrow HR and palpitation. - Partially \downarrow conversion of T_4 to T_3 . Dose: 10-40 mg t.d.s orally. c) <u>Antithyroid drugs:</u> Onset: Start their clinical effect after 2 weeks. Aim: To control thyrotoxicosis gradually till euthyroid state is reached.	• <u>Dose:</u> 160 μ Ci / gm of thyroid tissue. Improvement occurs by 8-12 weeks (2-3 months). If not, A second dose may be required.	• <u>Pre-operative preparation</u> a) <u>Routine Method:</u> (Medical treatment): 1- <u>Neomercazole</u> (<u>Carbimazole</u>): till euthyroid state is reached. 2- <u>Lugol's iodine</u> : 10-14 days immediately before surgery to \downarrow vascularity and friability of the gland. b) <u>Rapid preparation:</u> (β adrenergic blockers) Aim: Rapid control of C.V.S. manifestations. Action: Acts on Target organ and not on the thyroid itself. So, it must be continued for 1 week after the operation.	• <u>Pre-operative preparation</u> a) <u>Routine Method:</u> (Medical treatment): 1- <u>Neomercazole</u> (<u>Carbimazole</u>): till euthyroid state is reached. 2- <u>Lugol's iodine</u> : 10-14 days immediately before surgery to \downarrow vascularity and friability of the gland. b) <u>Rapid preparation:</u> (β adrenergic blockers) Aim: Rapid control of C.V.S. manifestations. Action: Acts on Target organ and not on the thyroid itself. So, it must be continued for 1 week after the operation.

DR. Ali Hassib.
 Prep. Yasmin Bassioumy.
 Re wr. Salah Yousuf.

<p>Preparations:</p> <p>1) <u>Lugol's iodine: (5% I + 10% KI)</u> Indication: Only in preoperative prep. Action: - ↓ TSH effect on thyroid gland. - ↓ Vascularity & friability of gland. - ↓ Iodine binding to tyrosine. - Storage of colloid within the acini. - It's action declines after 2 weeks. Dose: 10 drops t.d.s with milk or juice to mask the bitter taste.</p> <p>2) <u>Carbimazole: (the commonest drug used)</u> Action: - Blocks iodine binding to tyrosine. - ↓ Antibody titre. Dose: 10 mg t.d.s till euthyroid state is reached (2 months), then 5 mg t.d.s for 12-18 months. NB: 0.1 mg/day thyroxine with antithyroid drugs to: 1- decrease danger of producing iatrogenic hypothyroidism. 2- prevent goitrogenic effect (↑ size of gland) 3) <u>Propylthiouracil: (not used)</u> Action: - Blocks iodine binding to tyrosine. - Prevents peripheral conversion of T4 to active T3. 4) <u>Potassium Perchlorate: (not used)</u> Action: ↓ iodine trapping by the thyroid.</p>	<p><u>NB: Special problems in management:</u></p> <p><u>[1]-Thyrotoxicosis with pregnancy:</u> - Radioiodine is absolutely contraindicated: as destruction of fetal thyroid may result. - Antithyroid drugs at usual dose result in fetal hypothyroidism with goiter, may obstruct air ways minimal dose antithyroid drugs avoids this risk. - Surgery after short course of antithyroid and propranolol proved to be safe during 2nd trimester.</p> <p><u>[2]-Thyrotoxicosis in children:</u> - Radioiodine is contraindicated in children. - Surgery shows high recurrence rate. - Treated by <u>Antithyroid drugs</u> until late teens.</p> <p><u>[3]-Thyrocardiac patient:</u> - Thyroidectomy is ideal after control of the cardiac status. - In unfit patients: radioiodine is used with antithyroid drugs until the effect of radioiodine appears (6 weeks).</p> <p><u>[4]-Propotosis of recent onset:</u> - Not preferable to terminate the toxic status abruptly by surgery or radioiodine for fear of the risk of malignant exophthalmos. - So, antithyroid drugs are used until the proptosis becomes stable for 6 months, after which, thyroidectomy is permissible.</p>	<p><u>Operation:</u> The following precautions should be considered:-</p> <p>1) The incision should be wide with division of the pretracheal muscles, thus manipulation is minimal to avoid thyrotoxic crisis.</p> <p>2) The operation done is... <u>Subtotal thyroidectomy</u> Principle: - The main mass of the gland is removed leaving only a thin postmedial wedge of each lobe. - The amount of the gland left is 4-5 gm on each side. WHY? - To maintain an euthyroid state. - To preserve the parathyroids. - To protect the recurrent laryngeal nerve.</p> <p>3) Perfect haemostasis and free drainage to avoid thyrotoxic crisis.</p>
<p>Advantages</p> <p>1) Avoid surgical risks. 2) Avoid possible hazards of radioactive I.</p> <p>Disadvantages</p> <p>1) No way to predict which patient is liable to remission. 2) High relapse rate: 60% within 2 years from stoppage of treatment. 3) Further enlargement of the gland. 4) Adverse effects of drugs: Thiouracil & Carbimazole: GIT upsets, Rashes, Arthralgia and Agranulocytosis: (due to reversible bone marrow dep.) So, Bl. Picture must be done every 2 weeks.</p>	<p>1) Avoid surgical risks. 2) Avoid prolonged medical therapy.</p> <p>1) Thyroid insufficiency in up to 75-80% of cases after 10 years. So follow up is a must. 2) It is inadequate for treatment of 2ry toxic goiter (i.e. recurrence).</p>	<p>1) Rapid cure. 2) High rate of success.</p> <p>1) Morbidity and Mortality (negligible in experienced hands). 2) Recurrence rate (<5%). 3) Thyroid insufficiency (20-45%).</p>

	Renal stones +++++ مهم نظري	Ureteric stones +++ نظري	Vesical stones ++ محتمل نظري	Urethral stones + مستبعد نظري
Incidence	<ul style="list-style-type: none"> 10-20% of the population Males > females Middle age <p>HOWEVER, no age is immune</p>		✓ + Children	✓
Aetiology Metabolic (1-5)	<p>1-) Geography:- Mediterranean and deserts 2-) Climate and seasonal variations:- Hot countries</p> <p>3-) Water intake:- Quantity and quality (minerals and trace elements)</p> <p>4-) Diet:- *Oxalates= Spinach, tomato and mango *Uric acid: Purines= Meat and liver *Calcium= Milk</p> <p>5-) Metabolic and endocrinal factors:-</p> <p>[Ca oxalate and phosphate] [Uric acid] [Cystine and xanthine]</p> <p></p> <p>Hypercalcaemia Hyperuricaemia Hyperoxaluria</p> <p>6-) Infection:- urea splitting bacteria → alkaline urine</p> <p>7-) Stasis → Triple phosphate stone</p> <p>8-) Congenital anomalies</p>			
Types	<p>I-Ca oxalate and Ca phosphate stones (75%):-</p> <p>1-) Ca oxalate calculi:- Hard- radio-opaque- radiating specules</p> <p>2-) Ca phosphate calculi:- Yellow- radio-opaque -grow in alkaline urine- large in size e.g. Stag-horn stone</p> <p>II-Ammonium and magnesium phosphate stones:- yellow- radio-opaque- grow in alkaline urine- grow rapidly- friable</p> <p>III-Uric acid stones:- Yellow- smooth- hard - radio-lucent الوحيدة</p> <p>IV- Rare stones:- 1-) cystine 2-) Xanthine</p>			
<p>Edited By Ahmed Diaa AFTER Mostafa Mahmoud Hasanain</p>				
	C O M P L I C A T I O N S			
H Hematuria	✓	✓	✓	✓
I Infection (pyuria)	✓	✓	✓	✓
M Migration	✓	✓	✓	✓
O Obstruction				Retention of urine
M Malignancy	SCC	—	—	—

	Renal stones	Ureteric stones	Vesical stones	Urethral stone
Type of patient	Male- Middle aged	✓	✓+ Children	-
PAIN: Character Site Due to	<ul style="list-style-type: none"> • Dull aching • In the renal angle and referred to the ant. Renal point • Due to pelvic distension or stretch of the renal capsule 	<p> <u>I-Stones of upper 1/3:</u> • Colicky pain • In the loin referred to the groin, testicles or labia <u>II-As stones moves to the middle 1/3:</u> • Groin pain (iliac fossa D.D. append.) • Referred to the upper thigh, tip of penis or vulva <u>III-Stone impacted in the lower 1/3:</u> • Frequency and pain at the end of micturition • Severe • Referred to the tip of penis or vulva • Often associated with stranguary (= bladder tenesmus) </p>	<ul style="list-style-type: none"> • Varies from slight discomfort to severe agonizing pain • In supra-pubic region • Referred to tip of penis or vulva • Due to contraction of the bladder around the stone at the end of micturition • PAIN is more at daytime and aggravated by movement • STRANGURY: intense desire to micturate with passage of few drops of blood stained urine and a sense of incomplete evacuation • FREQUENCY 1st diurnal the diurnal and nocturnal(cystitis) & due to irritation of bladder mucosa especially trigone • DIFFICULTY IN MICTURITION: interruption of the stream or acute retention 	<ul style="list-style-type: none"> • Burning pain • During micturition • Interruption of the stream followed by retention of urine
Examination	During attack of renal or ureteric colic there is usually nausea and vomiting General: Fever is rarely present until there is UTI Abdominal exam: Moderate deep tenderness at the site of stone Often there is tenderness at post. Flank as well			
Investigations	I-Lab:- 1-)Urine analysis: **Crystals of same type that causes stones **Microscopic or gross hematuria **Pus cells 2-Urea and creatinine II-Radiological:- 1-)Plain X-ray"KUB":urinary calculi are 90% radio-opaque → → → III-Rarely needed : 1-)DMSA scan: if patient is sensitive to contrast 2-Ascending(retrograde) pyelography: for radiolucent calculi difficult to localize with other investigations			
			Renal Opposite T12, L1,2,3& may need lateral view- May be Staghorn Ureteric Opposite tip of T. process- on the sacral alae – On pelvic side wall Vesical Supra-pubic- midline Urethral Below symphysis 2-) IVU 3-)US	Exam usually can detect stone If metal bougie is passed شربة محمول CLICK is felt
			• Detection of obstruction • Helpful in radiolucent stones (filling defect) • Idea about renal functions	• Detect severity of obstruction(= back pressure changes) • Helpful in radiolucent stones • Difficult to view all the ureter

T R E A T M E N T

Renal stones		Ureteric stones	Vesical stones	Urethral stones
1-] During the attack	<ul style="list-style-type: none">HospitalizationAnalgesics (opiates)Antispasmodics e.g. AminophyllineAdequate hydrationAntibiotics in case of UTIFiltering of urine or micturating in a clean containerStone analysis (to detect its type)Planning for future therapy		2-] Conservative (Aim: تنزل لوحدها)	
3-] Measures to prevent recurrence	Advices:- 1-) High fluid intake especially in hot weather 2-) Certain precautions according to the type of stone, directed mainly to avoid certain diet 3-) Treatment of any metabolic disturbance e.g. GOUT and Hyperparathyroidism		Indications:- -Stone less than 5 mm diameter -No backpressure effect or infection -No distal obstruction Method:- 1-)The main item: High Fluid Intake 2-)During the attack of pain (the same previous measures)	
I N T E R V E N T I O N T R E A T M E N T				
R E N A L S T O N E S				
1-) E S W L				
Indications: <ul style="list-style-type: none">All stones < 2cmAs a part of combined therapy	Contraindications:- <u>UROLOGICAL:</u> <ul style="list-style-type: none">Large stone > 2cmAssociated pathologyDistal obstructionSolitary kidney: except with stenting of the ureter <u>NON-UROLOGICAL:</u> <ul style="list-style-type: none">Bleeding tendencyBone anomaly e.g. scoliosis or kyphosis (difficult localization)	<u>Advantages:</u> <ul style="list-style-type: none">PainlessSuitable for risky patientsSuccessful for Most U. stones except: radiolucent and hard stones	<u>Complications:</u> وهي نادرة 1. <ul style="list-style-type: none">Transient attack of hematuriaColicky uerteric painFeverFailure to disintegrate the stone	
2-) P C N L				
Indications: <ul style="list-style-type: none">If ESWL is CI or failedAs a part of combined therapy	<u>Contraindications:</u> <ul style="list-style-type: none">Absolute: Bleeding tendencyRelative:-Pregnancy-Bone anomalies-Cong. anomalies	<u>Advantages:</u> <ul style="list-style-type: none">Small endoscopic woundMild postoperative painShort hospital stay	<u>Complications:</u> <ul style="list-style-type: none">Rarely injury to:<ul style="list-style-type: none">*Duodenum*Spleen*Colon*PleuraComp. related to stone removal	<u>Method:</u> <ol style="list-style-type: none">Under fluoroscopic or US guidance, pass a guide wire through the flank to reach the renal pelvisDilate the trackPass a nephroscope with a sheath 24-30 frenchExtract the stone through the sheathlarge stone can be disintegrated before extraction by: US, Laser, Compressed air, Electrohydraulic waves "Direct contact lithotripsy"

TB	Lymphadenitis	Enteritis	Peritonitis	Renal	Bone	Joint
(1) <u>incidence</u> precipit. Fact = poor general resistance	** **	** **	** **	Young adult **	Children **	** **
(2) <u>Etiology</u> A. Organism B. Route	** Blood or lymph-	** -1ry: ingestion -2ry: blood born or swallowed sputum	** -Local +++ Blood born- (2ry 100%)	** Ascending infect. ((from bladder) Blood born	** Local- Blood+++	** -Local (from adjac. Bone) Blood-
(3) <u>Pathology</u> A. 1ry or 2ry B. types C. Tubercle (revealed by biopsy)	Both Lymph & blood born	Both Ulcerative (2ry) Hyperplastic (1ry)	2ry 3 ACE Ascitic Adhesive Acute millary Caseous encysted	2ry Primary & secondary + مخاطب و غرائب Sterile acid pyuria Ulcerocavernous TB Auto-nephrectomy	2ry -Encysted(brodie's abscess) Infiltrating- -Atrophic(sicca without caseation) Hypertrophic (الوحيد)with new bone formation	2ry Synovial or Osseous ((1ry affection)

TB	Lymphadenitis	Enteritis	Peritonitis	Renal	Bone	Joint
(4) <u>CUP</u>						
A. Type of Pt = incidence	**	**	**	**	**	**
B. General (TB Toxaemia)	**	**	**	**	**	**
C. Specific	Lymph \ Blood	Diarrhea-Colic- Loss of wt- + mass in (Hyperblastic)	Eg: Ascitic :Doughy ,Swelling Rolled omentum <u>Encysted:</u> Local cyst , Swelling	Frequency -Renal (pain, swelling) Vesical (pain, strangulate, tenderness)	Pain Spasm, wasting, swelling, abscess, deformity (SWSAD)	Pain SWSAD
D.complications						
- cold abscess	**	**			**	**
-Sinus	**	**	-----	-----	**	**
dissemination-	**	**			**	**
plus	Calcification 2ry infection Collar&stud	Perforation> fistula Stricture> IO		Destruction of kidney	Pathological fracture	Deformity- -Pathological dislocation- Paraplegia- Ankylosis-

D. tt of complications	Abscess Sinus	IO Internal fistula	Maybe for IO	-----	**	Paraplegia (local rest)
(7) <u>Different diagnosis:</u>	Other causes of: lymphadenopathy swellings at this site	----- Other causes of: IBD Mass Rt iliac fossa	----- Ascitic: other causes of ascites Encysted: Other causes of Abd cysts Adhesive: Other causes of IO	Other causes of: Frequency Pain in loin Mass	Chronic non specific osteomyelitis Iry & metastatic bone disease	Pott's disease of the spine (Diseased hip joint).. flexion deformity dt spasm of psoas ms

Many thanks to Dr : Aly Hassib




presented by: Amira Hassan



Nahla El Sedik

4



RENAL NEOPLASMS

	Renal Cell Carcinoma (Hypernephroma)	Wilms' Tumor (Nephroblastoma)
Introduction:	This is the commonest renal parenchymal tumor representing 75 % of all renal neoplasms.	This tumor arises from embryonic nephrogenic tissue and contains epithelial and connective tissue elements.
Incidence:	<ul style="list-style-type: none"> ✓ <u>SEX</u>: The male to female ratio is 2:1. ✓ <u>AGE</u>: The majority of patients are between 50 and 70 yrs. ✓ <u>SIDE</u>: Bilateral tumors are either synchronous or metachronous and occur in 1-2% of cases. 	<ul style="list-style-type: none"> ✓ <u>SEX</u>: Male to female ratio is equal. ✓ <u>AGE</u>: Peak incidence is 3 to 4 years.
Etiology:	<ul style="list-style-type: none"> 1. Using tobacco products. 2. Von-Hippel Lindau disease. 3. Loss of short arm of chromosome 3 is a constant finding. 	
Pathology:	<p>A) GROSSLY:</p> <ol style="list-style-type: none"> 1. The tumor usually starts in one pole of the kidney (usually the upper). 2. The color of the tumor is yellow due to high lipid content. 3. Areas of hemorrhage and necrosis. 4. There is an apparent pseudocapsule surrounding the lesion. 5. The tumor infiltrates the renal pelvis early in the course of the disease. <p>B) MICROSCOPICALLY:</p> <ol style="list-style-type: none"> 1. Adenocarcinoma that arises from the cells of the proximal convoluted tubules. 2. <u>Types</u>: <ul style="list-style-type: none"> ✓ <u>Clear cell type (usually)</u>: is due to cholesterol crystals in cytoplasm dissolve during preparation. ✓ <u>Granular cell type</u>: is due to the increased mitochondria in the cytoplasm. 	<p>A) GROSSLY:</p> <ol style="list-style-type: none"> 1. The tumor appears as a solitary sharply demarcated, apparently encapsulated mass with areas of hemorrhage and necrosis. 2. Renal pelvic invasion is rare. 3. Renal vein invasion may occur. <p>B) MICROSCOPICALLY:</p> <p>The tumor contains both:</p> <ol style="list-style-type: none"> 1. Epithelial cells that may form primitive glomeruli or tubules. 2. Connective tissue elements as areas of cartilage, fat, smooth or striated muscles are present. <p>Degrees of differentiation of the tumor vary from:</p> <p>Favorable histology (FH) to Unfavorable histology (UH).</p>



Renal Cell Carcinoma (Hypernephroma)	Wilm's Tumor (Nephroblastoma)
<p>C) SPREAD:</p> <ul style="list-style-type: none"> ✓ Direct Spread: The neoplasm soon infiltrates a) The renal pelvis b) The Zuckercandle's (Gerota's) fascia c) May even infiltrate adjacent structures. ✓ Lymphatic Spread: i. To the L.Ns. along the hilum of the kidney ii. Then to the paraaortic L.Ns. and finally to the thoracic duct. ✓ Blood Spread: Hematogenous spread gives rise to metastases in the lungs, bones, and brain. Sometimes a malignant thrombus may be present in the renal vein or extends even to the IVC. 	
<p>D) STAGING:</p> <p><i>Robson staging system for renal cell carcinoma</i></p> <p><u>Stage I</u> Tumor is confined to the kidney.</p> <p><u>Stage II</u> Tumor is confined to Zuckercandle's (Gerota's) fascia and involves the perinephric fat.</p> <p><u>Stage III</u> Tumor involves renal vein and/or regional L.N. + Malignant thrombus.</p> <p><u>Stage IV</u> Distant metastases (Lungs, Liver or Bone).</p>	<p>D) STAGING:</p> <p><i>National Wilms' tumor staging system (1989)</i></p> <p>This is a post-nephrectomy staging on which further management will depend.</p> <p><u>Stage I</u> Tumor is limited to the kidney.</p> <p><u>Stage II</u> Tumor extends beyond the kidney but is completely excised.</p> <p><u>Stage III</u> Residual non hematogenous tumor confined to the abdomen.</p> <p><u>Stage IV</u> Hematogenous metastases.</p> <p><u>Stage V</u> Bilateral renal involvement at diagnosis.</p>
<p>E) PROGNOSIS:</p> <p>According to stage.</p>	<p>E) PROGNOSIS:</p> <p>BAD.</p>
<p>Clinical picture:</p> 	<p>Type of the patient</p> <p>1) Hematuria حُمَاة</p> <p>Microscopic hematuria occurs, while gross hematuria is not common and denotes bad prognosis.</p>

	Renal Cell Carcinoma (Hypernephroma)	Wilm's Tumor (Neuroblastoma)
40% a) Stretch of the renal capsule. b) Passage of blood clots causing ureteric colic. c) Infiltration of the adjacent lumbar nerves. 30% An irregular hard, renal swelling. <i>The classical triad of hematuria, pain and renal mass is present in only 10 % of patients and they indicate an advanced disease.</i> May be the first presentation e.g. pulmonary or skull deposit.		One third of patients present with vague abdominal pain.
		2) Pain ألم
		3) Renal Mass كتلة
		4) Metastases انتشار
	5) Non Specific Syndromes: As fever, night sweat, weight loss or anemia.	5) Hypertension 60% It results from encroachment on the blood supply producing renal ischemia and rennin production.
6) Secondary Varicocele: A rapidly enlarging varicocele that doesn't empty on elevation of the scrotum 7) Systemic Syndromes: a) Hypercalcemia occurs in 5 % of cases due to secretion of a parathormone like substance by the tumor or due to the presence of bone metastases. b) Erythrocytosis. c) Amyloidosis.		6) Associations: Anomalies associated with Wilm's tumor 1. Hemihypertrophy. 2. Genito-urinary anomalies. 3. Aniridia. 4. Increase in the incidence of Neurofibromatosis.
Differential Diagnosis:	1. Other causes of mass in Rt or Lt hypochondrium e.g. In Rt: Hepatic mass and retroperitoneal mass In Lt: spleen and retroperitoneal mass. 2. Other causes of renal mass e.g. hydronephrosis. 3. Other causes of hematuria e.g. BPH.	1. Neuroblastoma: irregular tumor, hard in consistency and may cross the midline. 2. Hydronephrosis. 3. Multicystic kidney. 4. Infantile type of polycystic kidney. 5. Ewing's sarcoma A) LABORATORY: - Complete blood count, liver and renal functions. - Urine catecholamines help to rule out neuroblastoma.
Investigations		

	Renal Cell Carcinoma (Hypernephroma) B) RADIOLOGICAL: (For both) 1- <u>Plain X-ray (KUB)</u> : may show mottled calcification. (In RCC) 2- <u>IVU</u> : a. Enlargement of the kidney. b. Elongation, displacement, compression or amputation of a calyx. c. Displacement of the renal pelvis. d. Assessment of function of the other kidney. 3- <u>Renal Sonography</u> : • Can differentiate solid from cystic masses. • Can detect liver metastases.	Wilm's Tumor (Nephroblastoma) 4- <u>CT scanning</u> : • It can detect involvement or surrounding structures or involvement of contralateral kidney. • It can detect lymph node enlargement and invasion of the renal vein or inferior vena cava. • It can follow the tumor response to chemo and radiotherapy. N.B. CT scan has Replaced <i>Angiography</i> 5- <u>Chest X-ray</u> : To detect pulmonary metastases.
Treatment 	A) OPERABLE PATIENTS: 1- <u>Unilateral</u> : Radical nephrectomy is performed. It entails removal of the kidney within its Gerota's fascia together with the ipsilateral adrenal gland. Ligate the vascular pedicle as the first step of the operation for 2 reasons: 1) Prevention of dissemination of malignant cells during manipulation of the tumor. 2) To be able to remove a malignant thrombus from the IVC, if present. N.B. Patients who have hypernephroma or Wilm's in a solitary kidney are treated by partial nephrectomy with 1-2cm safety margin. 2- <u>Bilateral</u> : Patients who have both hypernephroma or Wilm's { Radical nephrectomy in the more affected kidney. Partial nephrectomy in the less affected kidney. }	B) INOPERABLE PATIENTS: (Criteria of inoperability) 1) Extensive local spread e.g. fixity to the post. abdominal wall and infiltration to the neighboring structures. 2) Extensive lymphatic spread e.g. para-aortic L.N. 3) Blood spread. 4) Unfit patient. 1- <u>Resectable</u> : remaining tumor in nodes or other organs should be marked with surgical clips to facilitate direction of radiotherapy. 2- <u>Irresectable</u> : should be treated with chemotherapy and are re-explored; usually the tumor can be removed.
HANY SAGR HANY AHMAD RAMY ALKAWAJESY	Chemotherapy • Interferon (alpha and gamma) • Interleukin-2 OR Radiotherapy.	Scheme of Postoperative management If there is no residual focal lesion: adjuvant chemotherapy using actinomycin D, vincristin, adriamycin If there is residual focal lesion: combination of radiotherapy and chemotherapy is applied. N.B. ttt of associations e.g. genito-urinary anomalies.






	General	Fracture ribs	Pneumothorax	hemothorax	Open pneumothorax(sucking)	Flail chest
Introduction	✓	✓	✓	✓	✓	✓
Etiology						
• Blunt	✓	✓	✓	✓	✓	✓
• Penetrating	✓	✓	✓	✓	...
• Blast	✓	✓	✓	✓	✓	...
• Iatrogenic	✓	✓	✓
• Spontaneous	✓	✓(muscular violence)	✓	✓
Types						
• Closed or open	✓	• ✓ simple or flail	Simple Open tension	• ✓ According to source: Systemic pulmonary	✓
Associated injuries	✓المكتوب	• Thoracic • Extrathoracic eg: abdominal				
Dangers that may cause death:						
• Respiratory insufficiency Upper airway obust. Retention of secretions	✓	✓(pain, flail chest)	✓ • Open(mediastinal flutter, pendulum respiration) • Tension(mediastinal shift, ↓-ve intrapleural pressure)	✓	✓	✓(paradoxical mov., pendulum resp., mediastinal flutter)
• Circulatory insufficiency	✓	✓(mediastinal flutter)	✓(↓ intrathoracic negativity, flutter, kink)	✓(hge, ↓ intrathoracic negativity)	✓(flutter)	✓(mediastinal flutter)
Clinical picture:						
• سطمية	✓	✓	✓	✓	✓	✓
• History of trauma	✓	✓	✓	✓	✓	✓
• General (assoc.inj)	✓	✓	✓	✓(shock)	✓	✓
• Specific	✓	✓	✓	✓	✓	✓
• Associated inj. eg:abdominal	✓	✓	✓	✓	✓	✓
Investigations:						
• سطمية(two)	✓	✓	✓	✓	✓	✓
• Of the disease: Routine:hb, hematocrite, bl.gases CXR Needle aspiration Monitoring (specific)	✓	✓	✓	✓	✓	✓
For associated inj. Eg:abdominal	✓	✓	✓	✓	✓	✓
Treatment:						
• سطمية	✓	✓	✓	✓	✓	✓
First aid:ABCDE	✓	✓	✓	✓	✓	✓
Definitive	✓	✓	✓	✓	✓	✓
Indic.for thoracotomy	✓	✓
Of associated inj	✓	✓	✓	✓	✓	✓
Of complications	✓(empyema)


Bone Tumors

	Osteosarcoma (giant cell tumor)	Osteosarcoma	Ewing's Sarcoma
Introduction	<p>It occurs only in mature bone. Most commonly in the distal femur, proximal tibia, proximal humerus and distal radius, but other bones may be affected.</p>	<p>It is the most common 1ry tumor of bone. It may affect any bone but most commonly involves the long bone metaphysis, especially around the knee and at the proximal humerus.</p>	<p>It's a rare tumor arising from vascular endothelium in BM. It usually occurs in the diaphysis of a long bone and gives rise to marked periosteal reaction.</p>
Pathology	<p>★ <u>Site:</u> usually from epiphysis</p> <p>★ <u>Gross picture:</u></p> <ol style="list-style-type: none"> 1. The tumor has a fleshy red appearance. 2. It comes quite easily in pieces when it is curetted but it is difficult to be removed completely from surrounding bone. <p>★ <u>Microscopic picture:</u></p> <p>multinucleated giant cells scattered in a background of stromal spindle shaped cells which may be responsible for determining the aggressiveness of the tumor.</p> <p>★ <u>The grading system</u> (I, II, III) has been discarded & all giant CT are considered potentially malignant.</p> <p>★ <u>Staging:</u></p> <p>Tx 1ry tumor can't be assessed. T0 No evidence of 1ry tumor. T1 Tumor is confined within cortex. T2 Tumor invades beyond cortex.</p> <p>Nx L.N. can't be assessed. N0 No L.N. metastasis. N1 Regional L.N. metastasis.</p> <p>M0 No distant spread. M1 Evident of distant metastasis.</p>	<p>★ <u>Site:</u> is in the metaphyses of long bones, where it destroys & replaces normal bone.</p> <p>★ <u>The rule of 80:</u> 80% of in teenage, 80% in LL, 80% around the knee, 80% in the lower end of femur, and 80% in metaphysis.</p> <p>★ <u>Gross picture:</u></p> <ol style="list-style-type: none"> 1. The tumor rapidly spreads towards the shaft, but it respects the epiphyseal cartilage and hence does not invade epiphysis or the joint. 2. <u>There are 4 main pathological features:</u> <ol style="list-style-type: none"> a. Bone destruction. b. Tumor bone formation (<i>sun-ray appearance</i>). c. Reactive bone formation (<i>Codman's triangle</i>). d. Soft tissue infiltration. <p>3. <u>Types</u> * :</p> <ol style="list-style-type: none"> 1. <u>Osteolytic type:</u> Soft, fleshy & vascular with areas of hge and necrosis. 2. <u>Scroding type:</u> solid & contains bone. <p>★ <u>Microscopic picture:</u></p> <ol style="list-style-type: none"> 1. The cell of origin is the primitive osteoblasts. 2. It shows considerable variation: Some areas may have the characteristic spindle cells with a pink-staining osteoid matrix & others may contain cartilage cells or fibroblastic tissue with little or no osteoid. <p>★ <u>Staging:</u> (the same) Applicable to all variants except the parosteal osteosarcoma.</p>	<p>★ <u>Site:</u> In the diaphysis of a long bone.</p> <p>★ <u>Gross picture:</u></p> <p>The tumor arises in the medulla as a semisolid grayish white mass which spread longitudinally and transversely raising the periosteum with subperiosteal new bone formation in successive layers parallel to the shaft (<i>onion peel appearance in X-ray</i>).</p> <p>★ <u>Cut section:</u> Ill-defined grayish white mass with areas of hge and necrosis.</p> <p>★ <u>Microscopic picture:</u></p> <p>Small rounded cells arranged in rosette around the blood vessels.</p> <p>★ <u>Staging:</u> (the same)</p>



AF

	Osteoclastoma (giant cell tumor)	Osteosarcoma	Ewing's sarcoma
<p>Clinical Picture:</p> 	 <p>✓ Type of patient: usually is a young adult (20-40 years of age). ✓ Pain at the end of a long bone. ✓ Sometimes slight swelling. ✓ There may be a history of trauma and pathological fracture in 10 - 15% of cases. ✓ On examination:- 1. Swelling at the end of a long bone. 2. The consistency depends on the degree of thinning of the expanded cortex; it may be soft, firm or egg-shell crackling sensation. 3. The neighboring joint is often irritated.</p>	 <p>✓ Type of patient: The incidence is highest between the age of 10 and 20 years, but a second peak occurs after 50 years due to malignant changes in Paget's disease. ✓ Pain is usually the 1st symptom; it is constant, worst at night & gradually increases in severity. ✓ Sometimes, patient presents with a lump but pathological fracture is rare because the patient is bed ridden due to the pain. ✓ On examination: Local tenderness & in later cases there is a palpable mass and the overlying tissues may appear swollen and inflamed, the regional L.N. may be enlarged.</p>	 <p>✓ Type of patient: patient age ranges from 10-20 yrs., usually in a tubular bone & esp. in the tibia, fibula or clavicle. ✓ Pain and swelling are the chief presenting features. ✓ The lump is warm, tender, ill-defined & diaphyseal with intermittent or continuous pyrexia. ✓ Blood born metastasis may occur to other bones and in lungs. ✓ Lymphatic spread to regional lymph nodes is common.</p>
<p>Investigations</p> 	<p>1. X-ray: A radiolucent area situated eccentrically at the end of long bone & has the characteristic soap bubble appearance. Important is the presence of the <i>medullary plug</i> (operculum) at the junction of the shaft with the tumor. The lack of this plug may signify malignant osteoclastoma.</p>	<p>1. Laboratory: ↑ ESR & ↑ serum level of alkaline phosphatase. 2. X-ray:- a- Streaks of new bone formation, radiating outwards from the cortex (<i>sun-ray appearance</i>). b- Reactive new bone forms at the angles of periosteal elevation (<i>Codman's triangle</i>). c- Ghost of cortex: hazy osteolytic areas of cortex. Both (a) and (b) are typical of osteosarcoma. But they may occasionally be seen in other rapidly growing tumors.</p>	<p>1. Laboratory: ↑ ESR & leukocytosis. 2. X-ray: diaphyseal areas of bone destruction with new bone formation in layers along the shaft (<i>onion peel effect</i>). More often the tumor extends into surrounding soft tissues and may show sunray appearance and Codman's triangle.</p>

	Osteoclastoma (giant cell tumor)	Osteosarcoma	Ewing's Sarcoma
	<p>3. CT scans and MRI will reveal the extent of the tumor, both within the bone and beyond.</p> <p>4. Biopsy: (essential) the edge is ideal</p> <ul style="list-style-type: none"> - The center is necrotic - To obtain tumor with surrounding normal tissue to determine the microscopic extent of spread. <p>5. Arthroscopy may be helpful to establish affection of articular surface</p>	<p>3. CT scans and MRI. (Discuss)</p> <p>4. Biopsy. (Discuss)</p> <p>5. Chest X-ray to detect pulmonary metastasis.</p>	<p>3. CT scans and MRI. (Discuss)</p> <p>4. Biopsy. (Discuss)</p> <p>5. Bone scan may show multiple areas of activity in skeleton.</p>
Differential Diagnosis	<p>1. Tuberculosis osteomyelitis.</p> <p>2. Brown tumors of hyperpara.</p> <p>3. Chronic osteomyelitis.</p> <p>4. Other 1ry bone tumor e.g. osteosarcoma or osteoclastoma (discuss shortly) & Ewing's sarcoma.</p> <p>5. Metastatic bone tumor.</p>		<p>1. 2rys from neuroblastoma (both have the same MP) usually the patient is below 5 yrs of age.</p> <p>2. Reticulum cell sarcoma.</p> <p>3. Osteomyelitis.</p>
Treatment 	<p>✓ The simplest treatment is curettage and bone grafting, but recurrence is common.</p> <p>✓ The treatment of choice is Wide excision, with replacement by specially designed prosthesis or by bone grafts.</p> <p>✓ Amputation is indicated for tumors which recur with increasing evidence of malignancy.</p> <p>✓ Radiotherapy is reserved for surgically inaccessible tumors.</p> <p>✓ Excision of the affected bone e.g. osteoclastoma of fibula.</p>	<p>1) Local control of the disease is by either amputation or wide local excision and prosthetic replacement. <i>The level of amputation should be proximal to the joints above the tumour, e.g. osteosarcoma at the tibia is treated by an above knee amputation.</i></p> <p>2) Cytotoxic drugs have improved the prognosis.</p>	<p>✓ The prognosis is usually poor.</p> <p>✓ The best results are achieved by combination of</p> <ul style="list-style-type: none"> - Chemotherapy. - Radiotherapy. - Surgery. <p>✓ Then a further course of chemotherapy for 1 year is given.</p>

RAMY ALKONATESY

SPLENOMEGALY & HEPATOMEGALY



A.H.		SPLENOMEGALY	HEPATOSPLENOMEGALY	HEPATOMEGALY
CONGENITAL		1. Cysts of spleen.		1. Reidel's lobe 2. Polycystic disease
INFECTIONS	BACTERIAL	1. Paratyphoid. 2. Typhus. 3. T.B. 4. Anthrax.	1. Typhoid. 2. Brucellosis. 3. Syphilis. 4. Abscess.	1. Leptospirosis. 2. Pyogenic cholangitis. 3. Pyelophlebitis. 4. portal pyemia.
	VIRAL	1. Psittacosis.	1. Infectious mononucleosis.	1. Viral hepatitis 2. CMV, 3. Herpes simplex
	PARASITIC	1. Malaria.	1. Bilharziasis. 2. Hydatid cyst. 3. Kala azar.	1. Amebic hepatitis . 2. Toxoplasmosis
BLOOD		1. Thrombocytopenia. 2. Thalassemia major.	1. Leukemias. 2. Hemolytic anemias. 3. Polycythemia vera. 4. Myelofibrosis.	1. Megaloblastic anemia.
METABOLIC		1. Gaucher's disease. 2. Porphyria. 3. Rickets.	Amyloidosis.	1. Wilson's disease. 2. Hemochromatosis . 3. Fatty liver e.g. Reye's, DM 4. Lipid storage. 5. Glycogen storage (von Geirkes).
CIRCULATORY		Portal vein occlusion: 1. Neoplastic 2. Thrombophlebitis	Portal HTN.	Chronic venous congestion: CHF, Pericardial effusion Constrictive pericarditis
TUMOURS	BENIGN	lymphangioma	Hemangioma.	
	MALIGNANT	1. Fibrosarcoma. 2. Waldenstrom macroglobinemia.	Malignant lymphoma.	1. Metastasis. 2. Hepatoma. 3. Holangioma.
COLLAGEN DISEASES			1. Felty's \$ 2. Still's disease 3. Sarcoidosis	

Large spleen:

1. Portal HTN (Bilharziasis)
2. Thalassemia major
3. Chronic myeloid leukemia
4. Splenic sarcoma
5. Gaucher's disease
6. Kala azar
7. Chronic malaria.
8. myeloproliferative disorders (PRV, Myelofibrosis)

Enlarged tender liver:

1. Congestion (HF, Constrictive pericarditis)
2. Infection (Amebic hepatitis & abscess, viral hepatitis, pyogenic abscess)
3. Malignancy

For every problem under the sun, There is the SYSTEM or there is none

Many Thanks To Dr. Aly Hassib .
DINA HASSAN (D.H.)

	General	Spleen	Liver	Intestine	Kidney
Introduction	✓	commonest	2nd		
Etiology:					
1. blunt	✓	✓	✓	✓	✓
2. penetrating	✓	✓	✓	✓	✓
3. blast	✓	✓	✓	✓	✓
4. iatrogenic	-----	✓	✓	✓	✓
5. spontaneous	-----	✓	✓	✓	✓
Pathology	-----	✓	✓	-----	Early- delayed
Complication	-----	✓	✓	-----	delayed
C/P:		Fatal - delayed - classic			
1. History of trauma	✓	✓	✓	✓	✓
2. General:					
a. Shock					
- hypovolemic	✓	✓	✓	✓	✓
- septic	✓	No	No	✓	✓
b. Associated	✓	✓	✓	✓	✓
3. Local					
- picture of wound	✓	✓	✓	✓	✓
- picture of inter. He	✓	✓	✓	✓	✓
Or					
Peritonitis	✓	No	No	✓	✓
4. Specific C/P	✓	✓	✓	-----	✓
Investigation:					
اسطنبة					
	Indication :				
1. Hb & Hematocrit	✓	✓	✓	✓	✓
2. Plain x-ray	✓	✓	✓	✓ Air under diaphragm	✓
3. U/S	✓	✓	✓	✓	✓
4. C.T	✓	✓	✓	✓	✓
5. DPL	✓	✓	✓	✓	IVP
6. Diagnostic laproscopy	✓	✓	✓	✓	DMSA
7. Inv for possible associated injury eg. Chest x-ray	✓	✓	✓	✓	✓
III:	Indication :				
1. A B C D E	✓	✓	✓	✓	✓
2. Antibiotic	✓	✓	✓	✓	✓
3. Incision	✓	✓	✓	✓	✓ SSOAP
4. Exploration	✓	✓	✓	✓	✓
5. Specific ttt	✓	✓	✓	✓	✓
6. TTT of associated injuries	✓	✓	✓	✓	✓

GENERAL I.O.

★ Definition.

★ Classification.

★ Etiology. →

★ Pathology. →

★ Clinical Picture:

- Symptoms ☐
- Examination ☐
 - General
 - Abd. Inspection
 - Abd. Palpation
 - Auscultation
 - Rectal exam.

- Of strangulation ☐


- Of complications.

★ Investigations:

- Laboratory.
- Plain X-ray.
- U/S.
- Double enema.
- Br. Enema.

★ III:

- Pre-operative.
- Conservative.
- Operative.



#

	INTUSSUSCEPTION	VOLVULUS
Pathology <ul style="list-style-type: none"> Proximally Distally At site Pathology of strangulation General lethal effects 	<p>★ At site: The ileocecal valve is invaginated into the cecum and may travel along the whole colon and may protrude through the rectum and anus.</p> <p>○ An intussusception consists of:</p> <ol style="list-style-type: none"> 1) Intussusceptum: Consists of an entering layer and a returning layer. 2) Intussusceptient: Is the outer ensheathing layer. 3) Apex: junction between the entering and the returning layer. It is always constant at the ileocecal valve. 4) Neck: Is the junction between the ensheathing & returning layer. <p>○ The mesentery containing the blood vessels maybe compressed between the entering and the returning layers → ischemia and gangrene.</p> <p>★ Pathology of strangulation. (Discuss)</p> <p>★ General lethal effects. (Discuss)</p>	<p>★ At site: The upper loop usually falls in front of the lower in anticlockwise direction resulting in a combination of obstruction together with occlusion of the main vessels at the base of the involved mesentery.</p> <p>○ It is, therefore, a form of closed loop obstruction & of strangulation obstruction.</p> <ol style="list-style-type: none"> 1. Closed loop obstruction: closed loop is formed within which the pressure rises rapidly, with high risk of gangrene and perforation. 2. Strangulation obstruction: Interference with the blood supply occur if it rotates more than 1.5 turn → strangulation and perforation follows with rapid fatal peritonitis. <p>★ Pathology of strangulation. (Discuss)</p> <p>★ General lethal effects. (Discuss)</p> <p>★ Type of patient: sigmoid volvulus is common in elderly constipated male.</p> <p>★ Symptoms:</p> <ol style="list-style-type: none"> 1. Pain: colicky and caused by the hyperperistalsis. 2. Vomiting is late. 3. Distension is huge & peripheral. 4. Absolute constipation is early. <p>★ Signs:</p> <ol style="list-style-type: none"> 1. General exam.: showed Hypovolemic & septic shock. 2. Abdominal exam.: <ul style="list-style-type: none"> ○ Inspection: huge distention. ○ Auscultation: Accentuated intestinal sounds. 3. Rectal exam.: Empty rectum, blood may be found on tip of examining finger. <p>★ Picture of strangulation. (Discuss)</p> <p>★ Picture of complications. (Discuss)</p>
Clinical picture <ul style="list-style-type: none"> Type of pt. Symptoms Signs Picture of strangulation Picture of complications 	<p>★ Type of patient: The common age is between 3 and 12 months. Male: Female = 2:1</p> <p>★ Symptoms: (ألم البطن)</p> <ol style="list-style-type: none"> 1. Pain: Attacks of colicky pain denoted by screaming, drawing the legs up to the abdomen, and pallor. The attacks alternate with intervals of apparent wellbeing. 2. Vomiting. 3. Distension. 4. Passage of bloody mucus (that looks like red current jelly) per rectum. <p>★ Signs: (Examination)</p> <ol style="list-style-type: none"> 1. General exam.: showed Hypovolemic & septic shock. 2. Abdominal exam.: <ul style="list-style-type: none"> ○ Inspection: Distention is not an early feature. ○ Palpation: There is emptiness of the right iliac fossa (Sign de Dance). ○ Auscultation: Accentuated intestinal sounds. 3. Rectal exam.: reveals the bloody mucus and sometimes the head of the intussusception can be felt. <p>★ Picture of strangulation. (Discuss)</p> <p>★ Picture of complications. (Discuss)</p>	<p>★ Type of patient: sigmoid volvulus is common in elderly constipated male.</p> <p>★ Symptoms:</p> <ol style="list-style-type: none"> 1. Pain: colicky and caused by the hyperperistalsis. 2. Vomiting is late. 3. Distension is huge & peripheral. 4. Absolute constipation is early. <p>★ Signs:</p> <ol style="list-style-type: none"> 1. General exam.: showed Hypovolemic & septic shock. 2. Abdominal exam.: <ul style="list-style-type: none"> ○ Inspection: huge distention. ○ Auscultation: Accentuated intestinal sounds. 3. Rectal exam.: Empty rectum, blood may be found on tip of examining finger. <p>★ Picture of strangulation. (Discuss)</p> <p>★ Picture of complications. (Discuss)</p>

Investigations	INTUSSUSCEPTION	VOLVULUS
<p>a. Laboratory investigations:</p> <ol style="list-style-type: none"> 1- Blood picture. 2- Blood urea, Serum electrolytes. <p>b. Imaging investigations:</p> <ol style="list-style-type: none"> 1- <u>Plain X-ray abdomen</u> 2- <u>Ultrasound examination</u> 3- <u>Br enema</u>: (In cases of doubtful diagnosis). Shows a cylindrical filling defect at the area of intussusception and there is arrest of further progress of the contrast (<i>claw sign or meniscus sign, and coiled spring sign</i>). <p>c. Double enema test (Discuss)</p> <p>1) Gastroenteritis: There is fever, vomiting, diarrhea & shock is very late, no mass felt on abdominal or P-R examination.</p> <p>2) Rectal prolapse: No manifestations of intestinal obstruction, a finger cannot pass around the protruding mass.</p> <p>3) Henoch-Schönlein purpura: No manifestations of I.O. + purpuric eruptions.</p> <p>4) Other causes of I.O. in this age group e.g. Strangulated hernia.</p> <p>Preoperative preparations: (Discuss)</p> <p>Conservative treatment: (Hydrostatic reduction)</p> <p>★ <u>Indications:</u> Early cases that show no evidence of peritonitis.</p> <p>★ <u>Methods:</u></p> <p>1) Barium (or air) is run in per rectum and is followed radiologically under the screen. The diagnosis is established and the reduction, affected by the pressure of the barium column, is monitored until it is complete.</p> <p>2) This is confirmed by visualization of appendix and the terminal ileum.</p> <p>★ <u>Contraindication:</u> Late cases or presence of abdominal distention or rigidity.</p> <p>★ <u>Complications:</u> Reduction of a gangrenous intussusception or perforation of the bowel.</p> <p>Surgical treatment:</p> <p>★ <u>Indications:</u> When hydrostatic reduction fails or the condition is advanced from the start.</p> <p>★ <u>Method:</u></p> <ul style="list-style-type: none"> ○ At laparotomy, the head of the intussusception is squeezed backwards out of the containing colon. ○ The proximal ileum should never be pulled backwards as this may lead to intestinal tears. 	<p>a. Laboratory investigations:</p> <ol style="list-style-type: none"> 1- Blood picture. 2- Blood urea, Serum electrolytes. <p>b. Imaging investigations:</p> <ol style="list-style-type: none"> 1- <u>Plain X-ray abdomen</u> 2- <u>Ultrasound examination</u> 3- <u>Br enema</u>: Plain X-ray shows the hug gas-filled sigmoid loop that may look like the inner tube of car tire (<i>Omega loop</i>). <p>c. Double enema test (Discuss)</p> <p>1) Other causes of acute abdomen</p> <p>2) Other causes of I.O. in this age group e.g. sigmoid carcinoma.</p> <p>Preoperative preparations: (Discuss) including proper colonic preparation in case of elective operation.</p> <p>Conservative treatment:</p> <p>★ <u>Indication:</u> Early cases with no evidence of gangrene.</p> <p>★ <u>Method:</u> Rectal tube is passed through a sigmoidoscope to untwist the sigmoid loop. Success will lead to gush of gas and fluid stools. The tube is left in place for later elective resection of the long sigmoid after proper preparation.</p> <p>Emergency Surgery:</p> <p>★ <u>Indications:</u></p> <ol style="list-style-type: none"> 1) Failure of conservative. 2) Late presentation. <p>★ <u>Method:</u> (<i>Mention differences between viable & non viable gut</i>)</p> <ul style="list-style-type: none"> ○ Gangrenous sigmoid is resected, the proximal end is brought out to the skin as a terminal colostomy and the distal end is closed by sutures (Hartmann's procedure); for later elective anastomosis. ○ Viable sigmoid is untwisted and either fixed to the posterior abdominal wall or resected as above. <p>★ The presence of gangrene or irreducible intussusception is an indication for bowel resection and anastomosis.</p> <ul style="list-style-type: none"> ○ Important to check the whole bowel as there may be multiple intussusceptions. ○ <i>Mention differences between viable & non viable gut</i> 	<p>a. Laboratory investigations:</p> <ol style="list-style-type: none"> 1- Blood picture. 2- Blood urea, Serum electrolytes. <p>b. Imaging investigations:</p> <ol style="list-style-type: none"> 1- <u>Plain X-ray abdomen</u>: Plain X-ray shows the hug gas-filled sigmoid loop that may look like the inner tube of car tire (<i>Omega loop</i>). 2- <u>Ultrasound examination</u> 3- <u>Br enema</u>. <p>c. Double enema test (Discuss)</p> <p>1) Other causes of acute abdomen</p> <p>2) Other causes of I.O. in this age group e.g. sigmoid carcinoma.</p> <p>Preoperative preparations: (Discuss) including proper colonic preparation in case of elective operation.</p> <p>Conservative treatment:</p> <p>★ <u>Indication:</u> Early cases with no evidence of gangrene.</p> <p>★ <u>Method:</u> Rectal tube is passed through a sigmoidoscope to untwist the sigmoid loop. Success will lead to gush of gas and fluid stools. The tube is left in place for later elective resection of the long sigmoid after proper preparation.</p> <p>Emergency Surgery:</p> <p>★ <u>Indications:</u></p> <ol style="list-style-type: none"> 1) Failure of conservative. 2) Late presentation. <p>★ <u>Method:</u> (<i>Mention differences between viable & non viable gut</i>)</p> <ul style="list-style-type: none"> ○ Gangrenous sigmoid is resected, the proximal end is brought out to the skin as a terminal colostomy and the distal end is closed by sutures (Hartmann's procedure); for later elective anastomosis. ○ Viable sigmoid is untwisted and either fixed to the posterior abdominal wall or resected as above. <p>★ The presence of gangrene or irreducible intussusception is an indication for bowel resection and anastomosis.</p> <ul style="list-style-type: none"> ○ Important to check the whole bowel as there may be multiple intussusceptions. ○ <i>Mention differences between viable & non viable gut</i>
<p>D.D.</p> <p>1) Gastroenteritis: There is fever, vomiting, diarrhea & shock is very late, no mass felt on abdominal or P-R examination.</p> <p>2) Rectal prolapse: No manifestations of intestinal obstruction, a finger cannot pass around the protruding mass.</p> <p>3) Henoch-Schönlein purpura: No manifestations of I.O. + purpuric eruptions.</p> <p>4) Other causes of I.O. in this age group e.g. Strangulated hernia.</p> <p>Preoperative preparations: (Discuss)</p> <p>Conservative treatment: (Hydrostatic reduction)</p> <p>★ <u>Indications:</u> Early cases that show no evidence of peritonitis.</p> <p>★ <u>Methods:</u></p> <p>1) Barium (or air) is run in per rectum and is followed radiologically under the screen. The diagnosis is established and the reduction, affected by the pressure of the barium column, is monitored until it is complete.</p> <p>2) This is confirmed by visualization of appendix and the terminal ileum.</p> <p>★ <u>Contraindication:</u> Late cases or presence of abdominal distention or rigidity.</p> <p>★ <u>Complications:</u> Reduction of a gangrenous intussusception or perforation of the bowel.</p> <p>Surgical treatment:</p> <p>★ <u>Indications:</u> When hydrostatic reduction fails or the condition is advanced from the start.</p> <p>★ <u>Method:</u></p> <ul style="list-style-type: none"> ○ At laparotomy, the head of the intussusception is squeezed backwards out of the containing colon. ○ The proximal ileum should never be pulled backwards as this may lead to intestinal tears. 	<p>a. Laboratory investigations:</p> <ol style="list-style-type: none"> 1- Blood picture. 2- Blood urea, Serum electrolytes. <p>b. Imaging investigations:</p> <ol style="list-style-type: none"> 1- <u>Plain X-ray abdomen</u> 2- <u>Ultrasound examination</u> 3- <u>Br enema</u>: (In cases of doubtful diagnosis). Shows a cylindrical filling defect at the area of intussusception and there is arrest of further progress of the contrast (<i>claw sign or meniscus sign, and coiled spring sign</i>). <p>c. Double enema test (Discuss)</p> <p>1) Gastroenteritis: There is fever, vomiting, diarrhea & shock is very late, no mass felt on abdominal or P-R examination.</p> <p>2) Rectal prolapse: No manifestations of intestinal obstruction, a finger cannot pass around the protruding mass.</p> <p>3) Henoch-Schönlein purpura: No manifestations of I.O. + purpuric eruptions.</p> <p>4) Other causes of I.O. in this age group e.g. Strangulated hernia.</p> <p>Preoperative preparations: (Discuss)</p> <p>Conservative treatment: (Hydrostatic reduction)</p> <p>★ <u>Indications:</u> Early cases that show no evidence of peritonitis.</p> <p>★ <u>Methods:</u></p> <p>1) Barium (or air) is run in per rectum and is followed radiologically under the screen. The diagnosis is established and the reduction, affected by the pressure of the barium column, is monitored until it is complete.</p> <p>2) This is confirmed by visualization of appendix and the terminal ileum.</p> <p>★ <u>Contraindication:</u> Late cases or presence of abdominal distention or rigidity.</p> <p>★ <u>Complications:</u> Reduction of a gangrenous intussusception or perforation of the bowel.</p> <p>Surgical treatment:</p> <p>★ <u>Indications:</u> When hydrostatic reduction fails or the condition is advanced from the start.</p> <p>★ <u>Method:</u></p> <ul style="list-style-type: none"> ○ At laparotomy, the head of the intussusception is squeezed backwards out of the containing colon. ○ The proximal ileum should never be pulled backwards as this may lead to intestinal tears. 	<p>a. Laboratory investigations:</p> <ol style="list-style-type: none"> 1- Blood picture. 2- Blood urea, Serum electrolytes. <p>b. Imaging investigations:</p> <ol style="list-style-type: none"> 1- <u>Plain X-ray abdomen</u> 2- <u>Ultrasound examination</u> 3- <u>Br enema</u>: Plain X-ray shows the hug gas-filled sigmoid loop that may look like the inner tube of car tire (<i>Omega loop</i>). <p>c. Double enema test (Discuss)</p> <p>1) Other causes of acute abdomen</p> <p>2) Other causes of I.O. in this age group e.g. sigmoid carcinoma.</p> <p>Preoperative preparations: (Discuss) including proper colonic preparation in case of elective operation.</p> <p>Conservative treatment:</p> <p>★ <u>Indication:</u> Early cases with no evidence of gangrene.</p> <p>★ <u>Method:</u> Rectal tube is passed through a sigmoidoscope to untwist the sigmoid loop. Success will lead to gush of gas and fluid stools. The tube is left in place for later elective resection of the long sigmoid after proper preparation.</p> <p>Emergency Surgery:</p> <p>★ <u>Indications:</u></p> <ol style="list-style-type: none"> 1) Failure of conservative. 2) Late presentation. <p>★ <u>Method:</u> (<i>Mention differences between viable & non viable gut</i>)</p> <ul style="list-style-type: none"> ○ Gangrenous sigmoid is resected, the proximal end is brought out to the skin as a terminal colostomy and the distal end is closed by sutures (Hartmann's procedure); for later elective anastomosis. ○ Viable sigmoid is untwisted and either fixed to the posterior abdominal wall or resected as above. <p>★ The presence of gangrene or irreducible intussusception is an indication for bowel resection and anastomosis.</p> <ul style="list-style-type: none"> ○ Important to check the whole bowel as there may be multiple intussusceptions. ○ <i>Mention differences between viable & non viable gut</i>




	ADHESIVE INTESTINAL OBSTRUCTION	PARALYTIC ILEUS
Definition	Arrest of downward propulsion of intestinal contents 2ry to intraperitoneal adhesions which constitute the commonest causes of intestinal obstruction in developed countries.	Arrest of downward propulsion of intestinal contents due to loss of propulsive power of the bowel leading to functional obstruction.
Etiology	Result from:- ★ <u>Post-operative</u> : previous abdominal surgery. It is commoner when talc is used to lubricate the surgical gloves. ★ <u>Post-inflammatory</u> adhesions e.g. after septic or tuberculous peritonitis. ★ <u>Congenital bands</u> . ★ Adhesions develop at the sites of ischemia of the bowel or at the scars of the abdominal wall.	★ <u>Reflex inhibition of intestinal motility</u> following abdominal operations, spine fractures, and retroperitoneal hemorrhage. <i>This may be due to sympathetic over action.</i> ★ <u>Metabolic abnormalities</u> as hypokalemia, uremia, and diabetic ketoacidosis. ★ <u>Peritonitis</u> due to direct toxic effect on the nerve plexuses of the intestine. ★ <u>Drugs</u> as anticholinergics (e.g. propanthine)
Pathology ▪ Proximally ▪ Distally ▪ At site ▪ Pathology of strangulation ▪ General lethal effects	★ <u>Proximally</u> : Distension (gaseous & fluid), Hyperperistalsis. (Discuss) ★ <u>Distally</u> : Empty collapsed intestine. ★ <u>At site</u> : ○ <u>Adhesions</u> may be multiple or solitary & have a tendency for recurrence. ○ <u>Obstruction</u> occurs by kinking or by direct obstruction of a small intestinal loop. ○ <u>Strangulation</u> may result from compression of the loop's blood supply. A band may induce localized ischemic necrosis by direct pressure on the intestine → perforation & peritonitis. ★ <u>Pathology of strangulation</u> . (Discuss) ★ <u>General lethal effects</u> . (Discuss)	★ There is marked distension of the small and large intestines with gas and fluid. (Discuss) ★ <u>General lethal effects</u> : The patient suffers a severe loss of fluid & electrolytes through accumulation in the distended bowel & through vomiting.
Clinical picture ▪ Type of pt. ▪ Symptoms 1. Pain 2. Vomiting 3. Distension 4. Absolute constipation ▪ Signs 1. General 2. Abd. Inspec. 3. Abd. Palpation 4. Auscultation 5. Rectal exam ▪ Picture of strangulation ▪ Picture of complications	★ <u>Type of patient</u> : ★ <u>Symptoms</u> : 1. Pain: is absent and is replaced by sense discomfort and fullness. 2. Vomiting. 3. Abdominal distention. 4. Absolute constipation. ★ <u>Signs</u> : 1. <u>General exam.</u> : showed Hypovolemic shock. 2. <u>Abdominal exam.</u> : ○ Inspection: Abdominal distention. ○ Palpation: -ve. ○ Auscultation: <u>Accentuated intestinal sounds</u> . 3. <u>Rectal exam.</u> : Empty rectum. ★ <u>Picture of complications</u> . (Discuss)	★ <u>Type of patient</u> : ★ <u>Symptoms</u> : 1. Pain: is absent and is replaced by sense discomfort and fullness. 2. Vomiting. 3. Abdominal distention. 4. Absolute constipation. ★ <u>Signs</u> : 1. <u>General exam.</u> : showed Hypovolemic shock. 2. <u>Abdominal exam.</u> : ○ Inspection: Abdominal distention. ○ Palpation: -ve. ○ Auscultation: <u>Accentuated intestinal sounds</u> . 3. <u>Rectal exam.</u> : Empty rectum. ★ <u>Picture of complications</u> . (Discuss)

Investigations	ADHESIVE INTESTINAL OBSTRUCTION	PARALYTIC ILEUS
Investigations a. <u>Laboratory investigations:</u> 1. Blood picture. 2. Blood urea, Serum electrolytes. b. <u>Imaging investigations:</u> 1. <u>Plain X-ray abdomen</u> (discuss) 2. <u>Ultrasound examination</u> 3. <u>Br enema</u> (to detect colonic obstruction) c. <u>Double enema test</u> (Discuss) 1) Other causes of acute abdomen 2) Other causes of I.O. in this age group e.g. Strangulated hernia. D.D. Treatment • Preoperative • Conservative • Operative	a. <u>Laboratory investigations:</u> 1. Blood picture. 2. Blood urea, Serum electrolytes. b. <u>Imaging investigations:</u> 1. <u>Plain X-ray abdomen</u> (discuss) 2. <u>Ultrasound examination</u> 3. <u>Br enema</u> (to detect colonic obstruction) c. <u>Double enema test</u> (Discuss) 1) Other causes of acute abdomen 2) Other causes of I.O. in this age group e.g. Strangulated hernia. <u>Conservative treatment:</u> ★ <u>Indications:</u> Early cases with no evidence of strangulation. ★ <u>Method:</u> o Naso-gastric suction, and intravenous replacement of losses (drip and suck). o Close observation is necessary to judge success which is indicated by resolution of pain and distention, passage of flatus, and the retrieval of clear gastric aspirate. <u>Surgical treatment:</u> ★ <u>Indication:</u> 1) Conservative treatment fails. 2) If there is evidence of strangulation or gangrene. ★ <u>Method:</u> Adhesions are divided and the bowel viability is assessed and dealt with accordingly.	a. <u>Laboratory investigations:</u> 1. Blood picture. 2. Blood urea, Serum electrolytes. b. <u>Imaging investigations:</u> 1. <u>Plain X-ray abdomen:</u> Multiple gas fluid levels. The gaseous distention includes the whole small and large intestines. 2. <u>Ultrasound examination</u> Other causes of I.O. (كتب الجدول الثاني) A) <u>Preventive:</u> 1. Preoperative correction of biochemical disturbances. 2. Gentle handling of the intestine during surgery. 3. For major abdominal surgery, a naso-gastric tube is used to decompress the bowel postoperatively. B) <u>Curative</u> 1. <u>Preoperative preparations:</u> Drip and Suck are the essentials of treatment. (Discuss) 2. Correction of underlying metabolic abnormalities and hypoproteinemia 3. If a postoperative ileus is unduly prolonged for more than 3 days, one should think of peritonitis from a leaking intestinal anastomosis and of mechanical obstruction from early fibrinous adhesions. Both Conditions necessitate reoperation.

 RAMY E AHMAD
 ALKONATEESY








Anal Canal

	Piles	Fissure (شرخ)		Fistula (ناصور)	Perianal suppuration or abscess (خراج)		Rectal prolapse						
		Acute	Chronic		Primary	Secondary							
Etiology	<p>Primary or secondary</p> <p>A) Secondary piles are due to:</p> <ol style="list-style-type: none">1) Pregnancy.2) Rectal carcinoma. <p>N.B. Portal hypertension leads to anorectal varices.</p> <p>B) Primary piles:</p> <p>Occur without any organic cause.</p> <ol style="list-style-type: none">1) Weakness in the wall of vein (hereditary)2) Chronic constipation & straining.3) <u>Anatomical factors:</u>a) Long column of venous bl.b) The veins are constricted as they pass through the ms. coat.c) Lax submucosa.	<p><u>Longitudinal tear in the mucous membrane lining the anal canal as a result of:</u></p> <ol style="list-style-type: none">1) Hard stool.2) Foreign body.3) Crohn's disease.		<p>Factors that help chronicity:</p> <ol style="list-style-type: none">1) The anal glands act as a reservoir for infection.2) Internal opening for infection.3) Fecal material.	<p>Primary</p> <p>Infection of the anal glands.</p> <p>Abscesses are classified anatomically into:</p> <ol style="list-style-type: none">1) Perianal abscess (60%).2) Ischiorectal abscess (30%).3) Submucous abscess (5%).4) Pelvirectal abscess (5%).	<p>Secondary</p> <p>After Crohn's disease.</p>	<p>A) Partial prolapse:</p> <p>*Common in children due to:</p> <ol style="list-style-type: none">1) Loss of the curve of sacrum.2) Loss of weight.3) Straining. <p>*May occur in adults due to:</p> <ol style="list-style-type: none">1) Advanced cases of hemorrhoids.2) Rectal polyp. <p>B) Complete prolapse:</p> <p>The condition is commoner in elderly, particularly females.</p>						
Pathology	<p>- Each pile consists of a dilated vein with an artery, CT & covered with mucosa.</p> <p>- They are arranged at 3-7-11 o'clock</p> 	<p>*Site: 90% of fissures are posterior due to Y-shaped arrangement of the superficial part of external sphincter with absence of support posteriorly.</p> <table><tr><th>Simple superficial linear tear along axis of anal canal.</th><th>Deep</th></tr><tr><td>No fibrosis.</td><td>The edges are fibrosed & indurated.</td></tr><tr><td>Ms. of anal canal are spastic & not fibrosed (relax under anesthesia)</td><td>Organic fibrosis of the anal sphincter.</td></tr><tr><td>Pain leads to ms. spasm leading to more pain. Acute fissure usually doesn't heal spontaneously.</td><td>Associated sentinel pile at its lower end (due to infection & oedema)</td></tr></table>	Simple superficial linear tear along axis of anal canal.	Deep	No fibrosis.	The edges are fibrosed & indurated.	Ms. of anal canal are spastic & not fibrosed (relax under anesthesia)	Organic fibrosis of the anal sphincter.	Pain leads to ms. spasm leading to more pain. Acute fissure usually doesn't heal spontaneously.	Associated sentinel pile at its lower end (due to infection & oedema)	<p>Classification of anal fistulae:</p> <p>A) Goodsall's rule</p>  <p>B) Old classification:</p> <ol style="list-style-type: none">1) Low anal fistulae: The internal opening is below the anorectal ring.2) High anal fistulae: The internal opening is above the anorectal ring. <p>C) Recent classification:</p> <p>See tft</p>	<p>Infection of these glands leads to formation of intersphincteric abscess which may spread:</p> <ol style="list-style-type: none">1) Downwards: Perianal abscess2) Outwards: Ischiorectal abscess3) Inwards: Submucous abscess4) Upwards: High intersphincteric abscess 	<p>Theories:</p> <p>Many theories e.g. It is case of sliding hernia due to weakness of the pelvic floor ms.</p>
Simple superficial linear tear along axis of anal canal.	Deep												
No fibrosis.	The edges are fibrosed & indurated.												
Ms. of anal canal are spastic & not fibrosed (relax under anesthesia)	Organic fibrosis of the anal sphincter.												
Pain leads to ms. spasm leading to more pain. Acute fissure usually doesn't heal spontaneously.	Associated sentinel pile at its lower end (due to infection & oedema)												

C / P	Swelling	*1 st degree: No prolapse, pt. may present by bleeding only. Diagnosed only by proctoscopy. *2 nd degree: prolapse only during defecation, reduced spontaneously at the end of defecation. *3 rd degree: prolapse during defecation & the pt. has to manually reduce them. *4 th degree: permanent prolapse of piles.		—	Present	Prolapse with same degrees as piles.		
	Pain	Absent except with complications		—	Severe			
	Discharge	+ pruritis		Excessive	—	Mucous		
	Bleeding	Bright red not mixed with stools which occurs at first with defecation but later at anytime		Slight	—	Present		
	Others			History of an abscess.				
On examination		Constipation & reflex symptoms (dysuria & dysmenorrhea)						
		*P/R: 1) Piles are not felt but seen through proctoscopy. 2) Exclude rectal carcinoma.	Seen with severe spasm of anal sphincter. (Don't do PR). Felt fibrosed with sentinel pile below.			Length of prolapse Mucoanal corrugations Prolapse thickness	Partial < 5 cm Absent Mucosa only	Complete > 5 cm Often present Rectal ms. thickness can be felt.
Complications	1) Bleeding. 2) Strangulation of prolapsed pile. 3) Gangrene. 4) Ulceration & infection. 5) Sloughing of strangulated piles. 6) Fibrosis of thrombosed pile. 7) Portal pyemia.	Abscess, fistula, fibrous contracture of the anal sphincter leading to stricture formation.				As piles		
DD		*DD of an anal fissure: 1) Tuberculous ulcers & syphilis 2) Carcinoma. *DD of painful anal conditions: 1) Anal fissure. 2) Peri-anal suppuration. 3) Prolapsed strangulated piles. 4) Acute peri-anal hematoma. (External piles) 5) Carcinoma of the anus.				1) Prolapsing hemorrhoids. 2) Prolapsing polyp. 3) Prolapsing intussusceptions.		

Investigations of anal fistula: (الفحص)

- 1) Proctoscopy shows the internal opening of the fistula
- 2) Colonoscopy &/or barium enema: to exclude an underlying specific cause.
- 3) Fistulography.

Treatment	Treatment is either conservative, injection or surgical.	Mainly conservative	Mainly surgical	Recent classification:	Urgent surgery by drainage (don't wait for fluctuation) under general anesthesia is performed.	1) In children:
	<p>1) <u>Conservative treatment:</u> Indicated in mild 1ry or 2ry piles. It consists of light non irritant diet, laxative, astringent ointment or suppository e.g. haemamolis.</p> <p>2) <u>Injection sclerotherapy:</u> *Indicated in: Uncomplicated 1st or early 2nd degree. *Method of injection: 5-20% phenol in almond oil is injected submucous & extravasous opposite the pedicle of the pile 3-5 cc are injected & one pile each time is injected at one week interval. *Complications of injection: 1) Sloughing due to injection in mm. 2) Submucous abscess. 3) Stricture. 4) Secondary hge. 5) Pain due to injection too low. 3) <u>Rubber band ligation:</u> (Barron's). 4) <u>Cryosurgery.</u> 5) <u>Photocoagulation.</u> 6) <u>Surgical ttt:</u> *Indicated in: 3rd & 4th degree piles.</p>	<p>1) Relieve constipation by liquid paraffin & dieting. 2) Anesthetic ointment before defecation. (Must explain to the pt.) 3) Digital dilatation under anesthesia to relax the sphincter & prevent associated spasm. * If these measures fail, internal sphincterotomy is done under general anesthesia.</p>	<p>1) If the fissure is NOT very chronic: - A lateral internal sphincterotomy operation is successful. - The internal sphincter is divided at the 3 o'clock position. - Relief of the spasm of the internal sphincter will allow healing. - The operation can be done by the closed or open method.</p> <p>2) If the fissure is chronic with a sentinel pile: - The best procedure is to do fissurectomy & posterior internal sphincterotomy. - A triangular segment with its apex upward & including the fissure, anal papilla & sentinel pile is excised. - The operation should include internal sphincterotomy as well.</p>	<p>1)  Inter-sphincteric</p> <p>2)  Trans-sphincteric</p> <p>For 1 & 2 → Do fistulectomy or fistulotomy</p> <p>3)  Supra-sphincteric (2 staged operation)</p> <p>4)  Extra-sphincteric -Difficult -Colostomy & staged operation.</p>	<p>N.B. The pt. who had drainage of an anorectal abscess should be warned that he may develop a perianal fistula later on.</p>	<p>1) <u>In children:</u> - Surgery is rarely indicated. *Simple measures are helpful: - Improving the nutritional status. - Avoid squatting.</p> <p>2) <u>In adults:</u> A) <u>Mucosal prolapse:</u> As piles. B) <u>Complete prolapse:</u> Various surgical procedures include: i) <u>Rectopexy</u> (Charles Well's operation)  ii) <u>Other option:</u> Treated as hernia by repair of pelvic floor ms.</p>

<u>Pilonidal sinus:</u>

*Pilonidal sinus:

Etiology: 1) Congenital theory 2) Acquired theory: Hairs are drawn through the skin.
Clinical features: Discharge maybe bloody & external opening.
DD: Perianal fistula

Treatment: 1) Pilonidal abscess: initially treated by incision & drainage of pus.

*Hemorrhoids (Piles): In Greek, hem means blood while rhoos means flowing. In Latin, pile means a ball. Piles may be internal or external according to its relation to the anal orifice.

*External piles = Acute perianal hematoma.

1st aid management:

- Airway: should be maintained patent.
- Analgesic: should be strong as 50ml pethidine IV.
- Tetanus prophylaxis.
- Tape water or saline at room temp. poured on burnt area for 15 min. to decrease edema and relief pain.

Minor Burns:

- **Criteria:** less than 15% in adults less than 10% in children
- Treated as out patient.
- A Analgesics
- A Antibiotics (systemic)
- Dressing using proper local chemotherapeutic.

Major Burns: & most moderate (except superficial)

- **Criteria:** Major: more than 30%
- **Moderate:** Adults 15% - 30% & Children 10% - 30%
- Administered to burn unit.
- C Canula: wide bore I.V. canula is inserted rapidly
- C Catheter: Foley's urethral catheter is inserted to monitor urine output.

Resuscitative Fluid Therapy

Amount and rate of fluid replacement depend on the weight of the patient and % of total body surface area injured.

- Amount infused during the 1st 24 hours is 2ml/percent surface area burn/kg bodyweight. Half the amount calculated is given over the 1st 8 hours & the other half over the next 16 hours. Also half the amount calculated is given over the 2nd day.

Formulas:

Evan's Formula: 1st day: 1 ml/kg/% normal saline + 1 ml/kg/% colloid + 2000 ml glucose. (the daily caloric needs)
2nd day: 1/2 ml/kg/% normal saline + 0.5 ml/kg/% colloid + 2000 ml glucose

Modified Brook's Formula:
1st day: 2-3 ml/kg/% lactated Ringer + 2000 ml glucose.
2nd day: 1 ml/kg/% lactated Ringer + 0.5 ml/kg/% colloid + 2000 ml glucose

Parkland's Formula:
1st day: 4 ml/kg/% lactated Ringer: 1/2 amount over 1st 8 hours
1/4 amount over 2nd 8 hours
1/4 amount over 3rd 8 hours

Duration: It is essentially given for the first 48 hours.

Monitored by: 1- Regular check up of vital signs
2- Urine output should be 30-60 ml/hour
3- C.V.P

N.B.:

- Max. % calculated in all formulas is 50% to avoid serious over infusion.
- Blood administration can be started after 48 hours guided with hematocrit value
- Oral intake is avoided in 1st 48 hrs to avoid GIT complications.
- Extensive burned patients are liable to have serious catabolic status. I.V. hyperalimentation makes it easy to correct this problem and to support the patient nutritionally during this critical period.

Local Wound Care

- The aim is to avoid infection.
- Constricting eschars 2be released immediately.
- Urgent fasciotomy in deeper burns may be limb saving.
- Topical antimicrobials should be applied after cleaning & conservative debridement.

Wound is managed by one of the following:

1. Exposure method: requires isolation in aseptic atmosphere.
Advantages: a. more comfortable to patient.
b. avoid repeated change of dressings.
c. inhibits bacterial growth by dry air.

Indications: a. Burns of face, neck & perineum.
b. Burns involving one side of the trunk.

2. Bulky Occlusive dressing (the occlusive method):
Since it is usually very painful; dressing change can be done by Hubbard Tank or under anesthesia (especially in children).
- Both methods are equally effective
- All partial thickness burns should heal within 2-3 weeks
- Full thickness burns require closure by autogenous skin grafts.

Biological Dressings:

Indications:

- a. Autograft is not enough
- b. Local wound condition is not favorable.

Advantages: a. Wound will be less painful.
b. Minimize fluid and protein loss. c. Control infection.

Examples:

- a. Allografts (cadaveric)
- b. Xenograft Skin (Pig's)
- c. Amniotic membrane.

- Applied after removal of eschar and are changed every 3-4 days.
- Only temporary and permanent closure only by autograft skin.
- Artificial skin substitutes are used with promising results.

Surgery

Time: 3-5 days post-burn (b4 bacterial contamination)
Method: tangential excision of damaged dermis & grafting is done either immediate or late when healthy granulations appear.
Indications: Deep burns

Prognosis Of Burn

- **Burn related factor:**
 - a. Extent: rule of 9
 - b. Depth
 - c. Type of burn: high tension electric burn has the highest mortality rate
 - d. Site of burn
 - e. Infection
 - f. Associated injuries
- **Patient Related:**
 - a. Age: extremes have bad prognosis.
 - b. Coexistent cardiopulmonary diseases.
- **TTT Related Factor:**
 - ttt in special burn centers gives better prognosis.

Post-op Comp

Name of surgery	Haemorrhage	✓ Infection	Injury	Paralytic ileus \ DVT\AGD\Incisional hernia	Specific complication
Hernia	✓	✓ mesh	Vas vessels	✗	Recurrence <i>Chronic inguinodynia</i>
Hydrocele	✓	✓	Testis Epididymis	✗	Recurrence
Varicocele	✓	✓	Vas Testicular artery	✗	Recurrence
Circumcision	✓ *	✓	Glans urethra	✗	Over circumcision Under circumcision phimosis
Appendectomy	✓	✓ Potentially contaminate wound	cecum	Incisional hernia	
Splenectomy	✓	✓ Subphrenic abscess ✓ overwhelming	Tail of pancreas Gastric wall	Incisional Hernia Paralytic ileus	Portal vein thrombosis
cholecystectomy	✓ *	✓ Subphrenic abscess	CBD	Paralytic ileus	Post cholecystectomy syndrome
Colostomy	✓	✓	Marginal artery	Paralytic ileus	Retraction Prolapsed Stricture Internal herniation Skin excoriation
Piles	✓ Primary ✓ Reactionary ✓ secondary	rarely	Sphincteric anorectal ring	✗	Stricture Reflex retention of urine Severe pain
Varicose veins	✓	✓	Saphenous Nerve Femoral Vain		Recurrence
Thyroidectomy	✓	✓	Recurrent laryngeal External laryngeal Trachea(surgical emphysema) Parathyroid(vascular injury/remove)	✗	Postoperative crisis Recurrence Hypothyroidism Myxoedema <i>Keloid</i> Progressive exophthalmos
Radical mastectomy	✓	✓	Axillary vessels Cephalic Vain N. to serratus N. to latissimus	✗	Briedel's scar <i>Seroma</i>
Kidney operations	✓	✓	Peritoneum Colon/deudenum Subcostal N.	Incisional hernia	Urinary fistula
BPH	✓ Primary ✓ Reactionary (clot retention)	✓	Sphincters Pudendal N.	✗	T.U.R syndrome stricture
Amputation	✓	✓	—	✗	Phantom limb Redundant soft tissue Spur formation Stump neuroma Adherent scar

Dr. A. H.

40

<p>Tetanus</p> <p>Definition:</p> <ul style="list-style-type: none"> - Specific anaerobic infection - Mediated by neurotoxin - Lead to \rightarrow Nervous irritability \rightarrow Tetanic muscular. Contractions 	<p>Gas Gangrene</p> <p>Definition:</p> <ul style="list-style-type: none"> - acute spreading gangrene. - Associated \hat{e} gas formation & profound toxemia by anaerobic spore bearing Bacilli
<p>Aetiology</p> <p>Organism :</p> <p>Clostridium tetani gram +ve</p> <p>Anaerobic Bacilli \hat{e} terminal</p> <p>Spore (drumstick appearance)</p> <p>Source of Organism</p> <ul style="list-style-type: none"> - Organism: naturally present in intestine of Horses - Spores present in manured soil & dirt - Spores resist heating, dryness, Boiling for 5 minutes <p>Mode of Infection</p> <ol style="list-style-type: none"> 1) Wound: contaminated by soil 2) Umbilical stump: infected catgut, contaminated dressing and powder. <p>Toxins</p> <p>Neurotoxin</p> <p>Predisposing Factors</p> <ol style="list-style-type: none"> 1) Wounds contaminated by hoarse excreta 2) Presence of FB or associated pyog. infection 3) Wound \hat{e} low oxygen tension e.g deeply seated & lacerated wound 4) Wound low Blood supply e.g anemia, shock, Tight bandage Plaster of Paris 	<p>Aetiology</p> <p>Organism: fall into two groups</p> <ol style="list-style-type: none"> 1- Saccharolytic Cl welchii cl. Septicum 2- Proteolytic cl. histolyticum cl. sporagers. <p>Source of Organism</p> <p>Organism: normal inhabitant intestine of man & animal</p> <p>Spores: manured land e.g Field battle- farms</p> <p>Mode of Infection</p> <ol style="list-style-type: none"> 1- Wound: contamination of extensive wound in: <ul style="list-style-type: none"> - War injuries - terrorist- attack on civilized person <p>Toxins α Toxin :</p> <p>Others include: hyaluronidase, lipase hemolysin</p> <p>Predisposing Factor</p> <ol style="list-style-type: none"> 1- lacerated wound involve Bulky ms e.g: gluteus 2- presence of FB or dead tissue 3- ischemia of muscle due to tight bandage, cast, suture undertension injury main vessel 4- infection \hat{e} aerobic bacteria \rightarrow make field suitable for clostridia(anaerobic) 5- Elderly person \hat{e} above knee amputation & fecal incontinence.
<p>Pathology</p> <ul style="list-style-type: none"> - neurotoxin is an exotoxin produced locally & reach CNS via blood or motor nerves or both. <p>Once reach CNS fixed by motor cells & then can't be detected in blood or CNS.</p>	<p>Pathology: either local or systemic</p> <p>Local: 1- saccharolytic org cause:-</p> <p>Necrosis of ms due to thrombosis of blood vessels and haemolysis of blood.</p> <p>Ferment glycogen of dead ms: liberation of H_2, CO_2</p> <p>Liberated BI pigment stain ms Brick red colour.</p>

<ul style="list-style-type: none"> - antitoxin can neutralize neurotoxin just before it's localized in CNS. - lead to excitability of medulla & spinal cord from which mild stimulus cause violent spasm - Death from toxemia, exhaustion, resp. obst. 	<p>2- Proteolytic org. cause: Ferment ptn. of dead ms \rightarrow liberation of H_2S $+Fe \rightarrow Fe_2S_3 \rightarrow$ Greenish black discoloration</p> <p>Systemic</p> <ul style="list-style-type: none"> - Blood hemolysis lead to \rightarrow pallor, Tinge of jaundice - Degenerative changes in liver, kidney, suprarenal
<p>Clinical Picture</p> <p>IP : non immunized pt 24h- 15 day Imm - 11 day may weeks or months</p> <p>divided into stages:</p> <p>Toxemia : Δ temp. Δ pulse Pt. Being irritable, headache, rigor Pt- gets Hepatitis, myocarditis, gn</p> <p>Tonic : Pain, numbness, lock jaw, neck stiffness, bitter smile</p> <p>Clonic : - violent ms. Contraction reflex to minor stimulus e.g light - stage characterized by</p> <div data-bbox="47 963 670 1209"> <pre> graph TD A[Incomplete relaxation] --> B[Arched Back] A --> C[Sweating profuse] B --> D[Marked Tachycardia] B --> E[Spasm at intercostals ms diaphragm] C --> F[Garve Prognosis] C --> G[prolonged apnea] </pre> </div>	<p>Clinical Picture</p> <p>IP : varies from few hours to few days.</p> <p>Divided : general, local</p> <p>1) General Δ temp. Δ pulse pt is apprehensive shock, oliguria</p> <p>2) Local:- gas: pain, numbness, wound is swollen, crepitus Gangrene: - serosanguineous discharge. - muscle: Brick red</p> <div data-bbox="702 918 1308 1041"> </div> <ul style="list-style-type: none"> - overlying skin: greenish black - offensive odour
<p>Investigation</p> <p>Polymorphnuclear leucocytosis is present</p>	<p>Investigation</p> <p>Mainly, diagnosis based on clinical appearance at wound.</p>
<p>DD. tt msr</p> <ol style="list-style-type: none"> 1) trismus: due to local cause e.g T.M.J arthritis 2) Tetany carpo-pedalspasm 3) Meningitis: neck muscle 1st. 4) Strychnine poisoning: complete relaxation 5) Rabies \rightarrow history of a dog bite spasm on seeing or drinking water Mainly MS. Of Deglutition & resp. 	<p>DD: A } other cl. Infection</p> <ol style="list-style-type: none"> 1) simple contamination not significant infection 2) gas abscess: - noninvasive, no ms. involvement - ttt : incision, drainage 3) cl. cellulitis - SC, no ms involvement - edema, gas, skin discoloration 4) Localized cl. myositis: - non invasive - myositis
<p>Complication</p> <p>Due to toxemia</p> <ul style="list-style-type: none"> - myocarditis - GN - Hepatitis - Avulsion fracture in bone <p>Clinical Subtypes See notes AH</p>	<ol style="list-style-type: none"> 5) oedematous gangrene : - Highly fatal - Cl. oedematiens - No gas B } Non cl infection Mixture of gram -ve bacilli & gram +ve cocci C } c- Surgical emphysema: presence of gas under skin

Prevention	Prevention
<p>Every child should be actively immunized by tetanus toxoid and continuing booster injection every 7- 10 years</p> <p>Individual on wound exposure</p> <p>• Proper immunized ↓ give booster dose 0.5ml IM</p> <p>• Improper immunized Clean → at risk ↓ • TT one exposure • TIG • Ab</p> <p>Not previous immunized Clean → at risk ↓ • TT 3 doses 4-6 weeks interval • TIG 250U IM • Ab</p>	<p>All clostridia infection are preventable.</p> <ul style="list-style-type: none"> - adequate debridement of wound, clean, left open - antibiotics: penicillin - adequate circ. support avoid tissue hypoxia - Anti GG serum isn't used.
<p>Treatment:</p> <p>Intensive ttt should be started soon</p> <p>1- neutralize by antitoxin TIG 3.000- 6.000 unit IM give in proximal portion of wound or in vicinity of wound, repeated doses may be required .</p> <p>2- Excise & debride: after neutralize of Toxin, wound left open & H2O2 may needed</p> <p>3- the pt .</p> <ul style="list-style-type: none"> - Aqueous penicillin 20- 40 million unit/ day - Barbiturates should be cautiously used to avoid CRS failure (depression) - Curarization & mechanical ventilation. - Don't disturb pt by unnecessary movement & excitement . - Dark quite room <p>N.B: One attack of tetanus doesn't give life long immunity</p>	<p>Treatment</p> <p>1) Wound Management (Under GA)</p> <ul style="list-style-type: none"> - Dead Tissues & muscle are excised - Decompression of Tight fascial compartment - deep fascia, skin left open - daily exam & debridement is necessary - diverting colostomy in extensive perineal infections - diffuse myositis & complete loss of blood supply or when a decquate debridement leave useless → amputation <p>2) Hyper baric oxygen ▼ bacteria invasion & toxin production given for 1-2 hrs, repeated every 6-12 hrs.</p> <p>3) Fresh blood transfusion (early given)</p> <p>4) Antibiotics: penicillin 20-40 million unit/day clindamycin metronedazole can be used</p> <p>Prognosis Mortality rate 20%</p>

Many thanks to Dr/ Aly Hassib

• Mohamed Farouk
"Rico"

• Abd'ElRahman Youness



Before



After

نصحه!
متنهزش

SYSTEM